

F&B Divosan TC 86

VS8

Chlorinated detergent-disinfectant for hard water

Description

Divosan TC 86 is a low foaming, chlorinated alkaline detergent-disinfectant for CIP and spray washing use in the food, beverage and dairy industries.

Key properties

- Divosan TC 86 contains a blend of caustic alkali, sodium hypochlorite and organic sequestrants/threshold
 agents. This combination of ingredients provides excellent soil and stain removal properties for use in hard
 water conditions. It also provides a highly effective disinfectant action against all types of microorganisms
 including bacteria, yeasts, fungi, spores and viruses.
- Divosan TC 86 is recommended as a single-stage cleaner and disinfectant for use in a wide range of CIP and spray washing applications. It is used in brewing and beverage plants for cleaning fillers, sugar/syrup tanks, open vessels, yeast handling and filtration equipment. It can be used in bars and restaurants for beer line cleaning. It is also used in dairies for cleaning milk reception areas and in food processors for general CIP of equipment and drain cleaning.
- Divosan TC 86 is low foaming and is suitable for use with a variety of CIP and spray washing equipment.

Benefits

- Broad spectrum disinfectant action.
- Highly efficient in the removal of organic material including food soil, stains and off-flavours.
- Low foaming under conditions of high turbulence, improves cleaning efficiency.
- Free-rinsing, minimises water usage.
- Cost-effective for use in hard water.

Use instructions

Use Divosan TC 86 at concentrations between 2-4% w/w (1.6-3.2% v/v) depending on the type and degree of soiling. Always rinse thoroughly with potable water after use. All food products and packaging materials shall be removed or carefully protected prior to usage.

For crate washing application use the product at concentrations between 0.5%-1% w/w (0.42%-0.83% v/v; at least 200ppm active substance) at least at 40 $^{\circ}$ C - 60 $^{\circ}$ C.

A virucidal effect can be achieved when using product concentrations above at least 1.75% w/w (1.46% v/v). Also used as automatic dish washer. Recommended concentration 0.3%.

Automatic and manual application and dosing.







F&B Divosan TC 86

VS8

Technical data

Appearance: Clear yellow to green liquid

pH (1% solution at 20°C): 12.5
Relative density (g/cm³; 20°C): 1.2
Chemical Oxygen Demand [COD] (gO₂/kg): 20
Nitrogen Content [N] (g/kg): 0
Phosphorous Content [P] (g/kg): 1.4

Divosan TC 86 [% w/w]	Specific conductivity at 25°C [mS/cm]
0.5	4
1	7.8
2	14.9
3	21.8
4	28.6
5	34.4

The above data is typical of normal production and should not be taken as a specification.

Safe handling and storage information

Full guidance on the handling and disposal of this product is provided in a separate Safety Data Sheet; sds.diversey.com. Store in original closed containers away from extremes of temperature. Only for professional users / specialists. Use disinfectants safely. Always read the label and product information before use and follow local regulations and advisory.

Product compatibility

Divosan TC 86 when applied at the recommended concentration and temperature is suitable for use on the grades of stainless steel commonly found in the processed food industry. It is unsuitable for use on cuprous materials and on soft metals such as aluminium. Always rinse surfaces thoroughly after use (within 1 hour). In the event of uncertainty, it is advisable to evaluate individual materials before any prolonged use.

Test method

Reagents

0.1 N Hydrochloric or sulphuric acid, 0.1 N Sodium thiosulphate, Phenolphthalein indicator

Procedure

Add approximately 10 ml of 0.1 N Sodium thiosulphate to 10 ml of test solution, mix well and allow to stand for about 30 seconds. Add 2-3 drops of indicator solution and titrate with acid to a colourless end point.

Calculation

% w/w Divosan TC 86 = titre (ml) x 0.36 % v/v Divosan TC 86 = titre (ml) x 0.3

Microbiological data

- $\bullet \quad \text{EN1276: passed at 1.5\% dilution in hard water (300ppm as CaCO_3) and soil (0.3\% bovine albumin), 5 minutes contact time.}\\$
- EN1650 (Candida albicans): passed at 1% (v/v) dilution in hard water (300ppm as CaCO₃) and high soil (3 g/l bovine albumin), 15 minutes contact time.
- EN13697 (Bactericidal): passed at 4% dilution in hard water (300ppm as CaCO₃) and low soil (0.03% bovine albumin), 5 minutes contact time and at 1.5% dilution in hard water (300ppm as CaCO₃) and low soil (0.03% bovine albumin), 30 minutes contact time.

www.diversey.com

© 2024 Diversey, Inc. All Rights Reserved. 10/04/2024 en (F00113)